

Amendments to the Claims: This listing of claims will replace all prior versions, and listings, of claims in the application

Listing of Claims:

1. - 12. (Cancelled)

13. (New) A method to obtain an immunotherapeutic agent that contains cell wall fragments from a virulent *Mycobacterium tuberculosis*-complex (MTB-C) strain of cells comprising the sequential steps of

- a) culturing the cells for a period of at least three weeks and
- b) homogenizing the cells in the presence of a non-ionic surfactant to produce a homogenate comprising non-fragmented cells, cell wall fragments, and solubilized cell compounds.

14. (New) The method according to claim 13 wherein the cell culturing period ranges from 3 to 4 weeks.

15. (New) The method according to claim 13 wherein the non-ionic surfactant is selected from the group consisting of alkylphenol ethoxylates and ethoxylated sorbitan esters.

16. (New) The method according to claim 15 wherein the non-ionic surfactant is an octylphenol ethoxylate compound.

17. (New) The method according to claim 16 wherein the non-ionic surfactant is an octylphenol ethoxylate having 7-8 mol of ethylene oxide.

18. (New) The method according to claim 13 wherein the cells are homogenized in a buffered medium having a neutral pH.

19. (New) The method according to claim 18 wherein the medium is buffered with PBS buffer.

20. (New) An immunotherapeutic agent obtained by the method according to claim 13.

21. (New) The method according to claim 13 further comprising the steps of
 - c) centrifuging the homogenized cell mixture to separate the cell wall fragments from the non-fragmented cells and the solubilized cell compounds,
 - d) washing the cell wall fragments and further treating the cell wall fragments to inactivate any remaining virulent cells, and
 - e) lyophilizing the resulting immunotherapeutic agent.
22. (New) The method according to claim 21 wherein the cell culturing period ranges from 3 to 4 weeks.
23. (New) The method according to claim 21 wherein the non-ionic surfactant is selected from the group consisting of alkylphenol ethoxylates and ethoxylated sorbitan esters.
24. (New) The method according to claim 23 wherein the non-ionic surfactant is an octylphenol ethoxylate compound.
25. (New) The method according to claim 24 wherein the non-ionic surfactant is an octylphenol ethoxylate having 7-8 mol of ethylene oxide.
26. (New) The method according to claim 21 wherein the cells are homogenized in a buffered medium having a neutral pH.
27. (New) The method according to claim 26 wherein the medium is buffered with PBS buffer.
28. (New) An immunotherapeutic agent obtained by the method according to claim 21.
29. (New) A pharmaceutical composition comprising the immunotherapeutic agent of claim 13.
30. (New) The pharmaceutical composition according to claim 29 in form of liposomes.

31. (New) The pharmaceutical composition according to claim 30 wherein the liposomes comprise auxiliary lipids selected from neutral and/or negatively charged phospholipids, and sterols.
32. (New) The pharmaceutical composition according to claim 30 wherein the phospholipids are selected from phosphatidylcholine, phosphatidylserine, and phosphatidylinositol.
33. (New) The pharmaceutical composition according to claim 30 wherein the sterols are selected from cholesterol and biliar salts.
34. (New) The pharmaceutical composition according to claim 30, further comprising vitamin E.
35. (New) A method for the combined treatment of tuberculosis comprising administering the immunotherapeutic agent of claim 20 in combination with at least one drug suitable for the treatment of tuberculosis.
36. (New) The method of claim 35 wherein the combined therapy is sequential or simultaneous.
37. (New) The method of claim 35, wherein the drug is selected from the group consisting of isoniazid, rifampicin, and combinations thereof.
38. (New) A pharmaceutical composition comprising the immunotherapeutic agent of claim 28.
39. (New) The pharmaceutical composition according to claim 38 in form of liposomes.
40. (New) The pharmaceutical composition according to claim 39 wherein the liposomes comprise auxiliary lipids selected from neutral and/or negatively charged phospholipids, and sterols.
41. (New) The pharmaceutical composition according to claim 40 wherein the phospholipids are selected from phosphatidylcholine, phosphatidylserine, and phosphatidylinositol.

42. (New) The pharmaceutical composition according to claim 40 wherein the sterols are selected from cholesterol and biliar salts.
43. (New) The pharmaceutical composition according to claim 39, further comprising vitamin E.
44. (New) A method for the combined treatment of tuberculosis comprising administering the immunotherapeutic agent of claim 28 in combination with at least one drug suitable for the treatment of tuberculosis.
45. (New) The method of claim 44 wherein the combined therapy is sequential or simultaneous.
46. (New) The method of claim 44, wherein the drug is selected from the group consisting of isoniazid, rifampicin, and combinations thereof.